



Early Covid-19 Treatment Guidelines: A Practical Approach to Home-based Care for Healthy Families

World Council for Health • VRS 6 • Current as of January 1, 2022

About this Guide

This is an up-to-date guide to managing Covid-19 illness effectively at home. Covid should be treated early, with a combination of therapies, and treated aggressively to avoid the more serious consequences of the illness.

As Covid-19 is a new condition, this guide is informed by both emerging medical research as well as the clinical experience of international doctor-led groups; it will evolve as new evidence emerges.

The safe, established and patent-free medicines and supplements included in this guide may have differing availability around the world; therefore a choice is provided. Expensive new drugs (such as monoclonal antibody treatments) are not considered here.

About World Council for Health

The World Council for Health is an umbrella organization of multinational groups of doctors, scientists and medicine journalists and other civil society members advocating for **the right to good health**. We ensure healthcare transparency through common-sense education and advocacy, independently integrating evidence from different qualitative and quantitative sources.





“A positive attitude is essential to a quick recovery!”

Who is this guide for?

Anyone can get Covid-19, whether they've had a Covid-19 shot or not. The good news is that the illness is easily treated, and most people will recover within a few days or weeks. By treating your illness early, you will be helping to end the Covid-19 pandemic.

If you or a family member have had a positive test or have symptoms suggestive of Covid-19, here is a practical guide to help you recover quickly. This guide is designed for people who are generally healthy and not taking other daily medications.

This latest guide was reviewed by:


Dr. Tess Lawrie, MD, PhD

Dr. Peter A. McCullough, MD, MPH

Dr. Pierre Kory, M.D., M.P.A.

Dr. Naseeba Kathrada, MD

We envision a world where you are empowered to take control of your health. However, this booklet does not provide individual medical advice or prescribe treatment but is provided as an educational service for patients and their families to know what options are available and widely used for many conditions. Readers should consult the physician(s) of their choice for individual medical evaluation and recommendations for treatment tailored to their individual needs.





In this guide, we will answer the questions:

- What are the symptoms of Covid-19?
- How does the illness progress?
- What can you do to treat Covid-19 at home?
- When should you go to the hospital with Covid-19?

We aim to empower you with information on practical, accessible medicines and therapies that will help you beat Covid-19 and get you back to living your life.

This guidance will evolve as new information on how to treat this viral infection emerges, so please sign up to World Council for Health email updates to get the latest developments on treating this new infectious disease.

What are the symptoms of Covid-19?


Covid-19 illness is thought to be caused by a virus called SARS CoV-2 virus. People infected with the virus will show any combination of the following:

- Fever or chills
- Cough
- Shortness of breath
- Difficulty breathing
- Fatigue
- Headache
- Loss of taste or smell
- Sore throat
- Congestion or runny nose
- Nausea or vomiting
- Diarrhea

Source: <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>



Phases & symptoms



Covid-19 has 3 phases

Much has been learned about Covid-19 since it was first identified. Scientists and clinicians now recognize 3 overlapping parts of the illness.

1. The Viral Phase (Day 1 - 5)

During the viral phase of the illness, the SARS-CoV-2 virus replicates rapidly in the body. Someone in the viral phase of the illness can expect flu-like symptoms such as fever, joint and muscle pain, headache, sore throat, loss of appetite, loss of taste/smell, nausea, diarrhea, and weakness.

Covid-19 transmission is highest during the viral phase of the illness, so avoid contact with people during this phase of the illness. It's important to note that asymptomatic people (those who test positive without symptoms) are unlikely to transmit the virus.

2. The Inflammation Phase (Day 5 - 10)

During the inflammation phase, your body's immune system response has geared up to fight the infection. A person in this phase of Covid-19 may continue to experience flu-like symptoms. Still, inflammation in lung cells may lead to feelings of breathlessness, coughing, or difficulty breathing and lead to pneumonia. If you are having difficulty breathing, you should go to the hospital. Covid-19 transmission is still possible during this phase.

3. The Hyper-inflammation/Clotting Phase (Up to 30 days)

Without early treatment, sometimes the symptoms progress into what's known as the hyper-inflammation or clotting phase or severe Covid-19. At this time, an infected person may experience additional symptoms, including chest heaviness/pain, further breathing difficulties, and blood clotting.

Useful Equipment

Having a **thermometer** at home is useful to check whether you have a fever. A temperature over 37.5°C that makes a person feel unwell can be treated to relieve symptoms.

A **pulse oximeter** is a small piece of equipment clipped onto a finger to measure the oxygen saturation of the blood. It is non-invasive and, in the context of Covid-19 illness, the test results are used to determine whether a person needs to be admitted to the hospital and/or receive

additional oxygen support. If you have access to a pulse oximeter, a threshold of 94% indicates you are not getting enough oxygen and should go to the hospital.

A **nebuliser** is a device that turns a solution into a mist, which then gets breathed in through a mouth piece or mask. It can be useful to deliver solutions such as saline to help with phlegm. Some international doctors also recommend nebulised colloidal silver, which has antiviral properties, and sodium bicarbonate solutions for Covid-19.




If you have a positive Covid-19 test or mild Covid-like symptoms

The first thing you should do is let your doctor know. Stay home, rest, drink lots of water, and follow the at-home guidance below. Early treatment will prevent disease progression.

If you have difficulty breathing, shortness of breath, chest heaviness, or are experiencing a sudden worsening of your symptoms, go to the hospital.

What can I use to treat Covid-19 at home?

Early At Home Treatment Essentials For Covid-19 Diagnosis or Symptoms



Over the course of the coronavirus outbreak, much has been learned by doctors and scientists around the world, and we now know how to treat Covid-19 at home.

With early treatment, the likelihood of a quick recovery goes up dramatically. It's a good idea to be prepared with simple medicines and therapies before you or your family members get sick.

The spike protein is a major component of the virus. It is toxic and affects the body in several different ways, which is why it's important to use a combination of the therapies to fight coronavirus disease. Infected individuals should use a combination of therapies that:

- support the immune system
- target the virus (antivirals/antimicrobials)
- reduce inflammation in the body (anti-inflammatories)
- reduce the risk of blood clots (anticoagulants)
- provide symptomatic relief for headaches, fever, cough, etc.

Most recommended therapies and medicines play more than one of these roles in the fight against Covid-19. Depending on availability in your area, prepare to treat with an agent in each of these five categories to maximize the chance of a speedy recovery.

Following is a basic 'shopping list' of options. Not all of these medicines and supplements are necessary. Aim to include an antiviral, anti-inflammatory and anticoagulant (aspirin), as well as a selection of immune supportive supplements in your treatment plan. Check the patient information leaflets to make sure that they are suitable for you as an individual.



**“Heal in the sun.
Our amazing sun
can heal us.”**

1. Antiviral/Antimicrobial Medicines

Ivermectin

Ivermectin (IVM) has antiviral properties, anti-inflammatory, and immune-supportive properties. This means that Ivermectin can be used in all phases of Covid illness, from early asymptomatic infection to more severe illness, including for hospitalized Covid-19 patients.

For early treatment of SARS-CoV-2 infection at home, the recommended dose is 0.4mg to 0.6mg per kilogram (kg) of body weight per day. In addition to being influenced by body weight, dosing might also be affected by the severity of the coronavirus infection and the virulence of the variant.

Ivermectin is widely used as an anti-parasitic medicine and, as such, is available in some countries over-the-counter (OTC). It comes in 3mg, 6mg, and 12mg tablets. Some countries also have it available as a liquid suspension.

For non-hospitalized adults of average body weight (between 50 to 80kgs), the dose of ivermectin is approximately two to four 12mg tablets daily with food for five days. Longer treatment or higher doses may be necessary, but this should be discussed with a doctor.

Side effects include nausea, diarrhea, dizziness, and rash. They tend to be mild and usually resolve upon discontinuation. Ivermectin is not recommended for use in children under 5 or



pregnant women (particularly in the first trimester) without the advice of a doctor. These side effects can be with minimised of the use an antihistamine like loratadine (Claritin).

If you take blood-thinning medication, please consult a doctor before taking ivermectin, as ivermectin also has blood-thinning properties.

Mouthwashes and Nasal rinses

Because so much virus is carried in your mouth and nasal passages, it is useful to rinse them once or twice a day with a saline and bicarbonate mixture or an antimicrobial mouthwash. Rinsing can also provide symptom support for those with excess mucus and phlegm.

Hydroxychloroquine

Hydroxychloroquine (HCQ) is a widely used anti-malarial drug. Its antiviral properties make it a useful alternative to ivermectin in the early treatment of Covid-19 at standard doses (200mg every 12 hours for 7 days). Doctors sometimes prescribe hydroxychloroquine and ivermectin together.



Doxycycline

Doxycycline is an antimicrobial drug commonly used to treat acne and Lyme Disease and prevent malaria. As it also has antiviral properties, many covid care experts use doxycycline (100mg every 12 hours for 7 days) and ivermectin together in early infection.

For other antivirals that may be available in your setting, please see the list at the end of this guide.

Antibiotics (for symptoms related to lungs)

If you develop chest symptoms, your doctor may need to prescribe you an antibiotic to prevent or treat suspected pneumonia.



2. Anti-inflammatory medicines

Anti-inflammatory medicines include steroids (e.g., budesonide, prednisone, dexamethasone, methylprednisolone), non-steroidal anti-inflammatory medicines (e.g., ibuprofen and aspirin), and certain other well-established medicines (e.g., colchicine and ivermectin).

Ibuprofen

Ibuprofen is a non-steroidal anti-inflammatory drug that is used to treat pain by reducing inflammation in the body. It is available for general use in most countries and helps to reduce the inflammation associated with Covid-19 and to treat symptoms, such as headache.

N-acetylcysteine


Glutathione is a powerful antioxidant that is found in most cells of the body. Its role is to protect cells by neutralizing toxins. Glutathione often becomes depleted during periods of illness. N-acetylcysteine is a drug that breaks down toxic substances by replenishing glutathione. N-acetylcysteine also helps loosen thick mucus in the inflammatory phase of covid illness.

Anti-histamines

Anti-histamines are usually used to treat allergies. They work by blocking the action of histamine; a chemical released when you have an allergic reaction. Oral antihistamines such as loratadine and cetirizine, commonly used OTC medicines for hayfever and rash, may help prevent an allergic immune response (mast cell activation) to the spike protein.

Steroids

Steroids are available by prescription only. Your doctor may prescribe budesonide, prednisone, dexamethasone, and methylprednisolone to help reduce inflammation. These are usually reserved for severe or persistent infection and are given under careful supervision by your doctor.



3. Immune-supportive medicines and supplements

Many nutritional supplements like Vitamin D, Vitamin C, Zinc, Quercetin, Melatonin, Glutathione, and N-acetylcysteine have powerful anti-inflammatory, antioxidant and immune-balancing properties. These therapies are simple, generally safe, and accessible for home-based treatment, and they do not require a prescription.

Vitamin D

Vitamin D plays an important role in immunity. A deficiency in vitamin D has been correlated with more severe Covid-19 illness. Maintaining good levels of this vitamin is critical for Covid-19 prevention and treatment. The oily vitamin D capsule is preferable to the tablet form, but you may use whatever is available and affordable. Vitamin D is best absorbed with a fatty meal.

Vitamin D is produced by the skin when it is exposed to sunlight but is not adequate in most places. Getting daily sunshine is vital for both the prevention and treatment of Covid-19.

Vitamin C and Zinc

Similarly, Vitamin C and Zinc have antiviral, anti-inflammatory, and antioxidant properties and are used in various viral illnesses, including Covid-19. Vitamin C can be found in many fruits and vegetables, especially citrus fruits, or taken as a supplement. Zinc is available in foods like beef, chicken, and fish. In the context of Covid-19 infection, you should use vitamin C and zinc supplements to ensure adequate levels are achieved.


Quercetin

Quercetin is a plant pigment or flavonoid widely available in fruit and vegetables, most notably parsley, berries, onions, apples, and citrus fruit. Quercetin can be taken as a supplement to help support the immune system during an infection. If you're also taking ivermectin, take quercetin at a different time of day.

Melatonin

Melatonin is a hormone made in the pineal gland in the brain. Studies show that melatonin reduces cytokines (proteins associated with inflammation) and improves sleep quality which is vital during a viral illness. Cherry juice contains low levels of melatonin. When used as a supplement, melatonin is best taken before bed.





4. Anticoagulant medicines

A dangerous feature of severe Covid-19 is blood clotting. Anticoagulant medications are used to prevent blood clots from forming.

Aspirin

As a cheap and widely available medicine, daily aspirin is recommended for early Covid-19 infection to prevent this serious complication from occurring. Aspirin has been in use for a long time for various health conditions and is available as an over-the-counter medicine. Its role in Covid-19 is to inhibit platelet aggregation, which reduces the risk of blood clotting. It also reduces inflammation and treats headaches. Aspirin is not recommended for children under the age of 16 or pregnant women in the last three months of pregnancy.

Prescription medications: Enoxaparin, heparin, and rivaroxaban

Not to be used without doctor supervision. Rivaroxaban is an oral anti-clotting medication that your doctor may prescribe for use at home. Enoxaparin is used for the prevention of deep vein thrombosis and pulmonary embolism. Because of its action in preventing blood clots, it can help treat worsening Covid-19. Enoxaparin and heparin are given by injection. Heparin is usually reserved for hospital use.

5. Symptom support

Many over-the-counter medicines and home remedies can relieve symptoms and help a person feel more comfortable, including cough mixtures, antihistamines, acetaminophen (paracetamol, Tylenol), as well as nasal sprays, mouthwashes, and other mucus-reducing agents. Saline or sodium bicarbonate solutions are effective and easy to prepare. Alternatively, to get rid of mucus in your lungs, you can try inhaling some steam with a tea towel over your head. Ginger and honey can help to relieve coughing – try a ginger tea with hot water and fresh ginger.





Adult Dosing Chart for At-Home Treatment of Covid-19

World Council for Health • Current as of January 1, 2022

<i>Treatment</i>	<i>Dosage</i>
Immune Support	
Vitamin D Vitamin C Zinc Quercetin Melatonin	10,000IU for 2-3 weeks, then 4,000IU daily 2 – 5g then .5 – 1g per hour 50mg once daily 1 tablet twice daily 5 – 10mg daily
Antiviral	
Ivermectin Mouth/nasal wash	0.4–0.6 mg/kg body weight for 5 days Rinse 3 times daily
Anti-Inflammatory	
Ibuprofen N-Acetylcysteine Anti-histamines	400mg up to 3x's daily 1–2 tablets daily for 7 days Use as directed on package
Anticoagulant	
Aspirin	300–325mg daily (1 tablet)
Symptom Support	
Paracetamol/ acetaminophen Cough medicines	Use as directed on package Use as directed on package

Child Dosing Chart for At-Home Treatment of Covid-19

World Council for Health • Current as of January 1, 2022

<i>Treatment</i>	<i>Dosage</i>
Prevention	
Vitamin D	1000IU per 15kg body weight
Vitamin C	1000mg
Zinc	20mg
Elderberry chewable	100 - 150mg
Melatonin	1mg
Treatment	
Vitamin D	1000IU per 15kg body weight
Vitamin C	1000mg
Zinc	20mg
Elderberry chewable/syrup	100 - 150mg
Melatonin	1mg
N-acetylcysteine	500mg
Azithromycin	10mg/kg body weight
Black seed oil	40 - 80mg/kg body weight
<i>Choose one:</i>	
Quercetin	250 - 500mg 2x daily
Hydroxychloroquine	100mg once daily
Ivermectin	0.2mg/kg bodyweight

Note: For severe infection, your doctor may prescribe steroids.



Remember: Treat Early with a Variety of Therapies

Treat Covid-19 early with a combination of antiviral medicine, anti-inflammatory therapy, anticoagulant medicine, simple immune-supporting therapies, and other medicines that will make you more comfortable during infection.

The medications and therapies you choose will depend on availability in your area. Use a selection of what is affordable and available to you. You don't need to use all of them!

Rest often and stay positive

Remember that the vast majority of people recover from Covid-19. Treat early and get plenty of rest, sunshine, and fresh air to reduce your illness's length and severity. Drink plenty of water.

Do not be in a hurry to resume your usual activities, e.g., going to the gym and social engagements—doing so may set your recovery back. Rather, allow yourself a full two weeks to recuperate. Contact your doctor if you have any questions or concerns.

When do I need to go to the hospital with a Covid infection?

SARS-CoV-2 infections range in severity. Some people will have no symptoms, some will have mild symptoms, and some will have severe symptoms that require hospitalization. Whether you have a confirmed Covid-19 infection or not, don't delay seeking advice from a medical professional. Early treatment is key to prevent severe Covid-19.

Consult your doctor or health authority:

- if you are worried about your symptoms
- if you are unsure what to do
- for information on early treatment options

Adults should go to the hospital if they:

- are short of breath
- are light-headed (this can be a sign of decreased oxygen in your blood)
- are having difficulty breathing
- are experiencing chest heaviness or pain
- have bluish lips
- Persistent fever, or other symptoms for more than 7 days since onset

Children should go to the hospital if they:

- are not interested in eating or breastfeeding
- have a blue tinge to their lips
- are confused
- have shortness of breath or difficulty breathing

Source: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/media-resources/science-in-5/episode-36---safe-care-at-home>

Be prepared. Keep a supply of basic medicines and therapies at home for your family.

The coronavirus pandemic has had a huge impact on physical and mental health across the world. Still, as we learn more about the illness, we can better treat and manage it so that severe illness is increasingly rare. The most important things you can do are to be prepared, take care of your immune system, and treat early using a combination of therapies that attack covid from a variety of different angles.

And remember, it's important to:

1. Maintain a healthy immune system with good nutrition, fresh air, and simple supplements like Vitamin C, D, Zinc, and Quercetin
2. Use the medicines and supplements that are available to you

3. Be prepared! Keep a good supply of healthy food, supplements, and medicines so you will be ready if you become infected.
4. Treat Covid-19 early if you become symptomatic. Early treatment helps reduce the severity and duration of Covid-19 infection
5. Treat Covid-19 using a combination of therapies that address the complex nature of the illness:
 - Antiviral/antimicrobial therapies such as ivermectin
 - Anti-inflammatory medicines such as ibuprofen and N-acetylcysteine
 - Immune-balancing medicines and supplements such as vitamin D, vitamin C, zinc, quercetin, and melatonin
 - Anticoagulant medicines such as aspirin
 - Symptom support such as cough mixtures, acetaminophen, nasal sprays, and mouthwashes

Seek advice from your doctor if you are concerned about Covid-19 symptoms or for early treatment options

GO TO THE HOSPITAL if you are short of breath, have difficulty breathing, or are experiencing chest heaviness or pain

Take care to rest and resume normal activities slowly after a Covid-19 infection.

Additional Resources

Summary of the Medicines used to Treat Covid-19

This list of medicines comprises over-the-counter (OTC) and prescription medicine for you to share with your doctor. Be sure to read the patient information leaflet before using any OTC medicines to check whether they are suitable for you as an individual, especially if you are pregnant. Aspirin and ibuprofen are not suitable for pregnant women, particularly in the last trimester of pregnancy. If your symptoms persist beyond 7 days, you feel your condition is worsening, your condition deteriorates suddenly, or if you are short of breath, contact your doctor or go the emergency room.

If you need to go to see a doctor or go to the hospital, print this list of medicines out and take it with you.

Don't forget: Drink lots of water, take lots of rest, eat lots of fruit and vegetables, and get daily sunshine. Allow yourself at least two weeks to recover without strenuous activity. Do not expect to resume your usual exercise regimen and social engagements for at least two weeks. A positive attitude is essential!

Immune Support

Multivitamin	1 tablet once or twice daily
Vitamin D3	10,000IU for 2-3 weeks, then 4,000IU daily 2x Dose: Care home residents & people with highly melanated skin
Vitamin C (ascorbic acid)	2-5g, then .5-1g per hour
Zinc	50mg once or twice daily
Omega-3 fatty acids	2-4g daily
Vitamin A	1 tablet daily
Vitamin B complex	1 tablet daily
Quercetin	1 tablet twice daily
Melatonin	5-10mg daily
Lactoferrin	200mg twice daily
Black seed (Nigella sativa)	1 capsule daily
Neem	1 capsule once or twice daily, not for combination use with B-Complex supplements
Curcumin	1 capsule once or twice daily
Lugol's Iodine	25mg daily as solution or tablets

Antiviral/antimicrobial

Antimicrobial mouth wash	Various OTC preparations, see below.
Antimicrobial nasal spray/rinse	Various OTC preparations, see below.
Azythromycin*	500mg daily for 5 days
Colloidal silver	5mls nebulised daily
Doxycycline*	100mg tablet twice daily for 7 days

Antiviral/antimicrobial As prescribed

Hydroxychloroquine*	200mg tablet twice daily for 5 to 7 days
Ivermectin*	0.4–0.6mg/kg body weight for 5 days
Nitazoxanide	500mg 2x daily after a meal for 5 days or until recovered

Anticoagulant As prescribed

Aspirin	300–325mg (1 tablet) daily
Rivaroxaban*	As prescribed
Enoxaparin/Low molecular weight heparin*	As prescribed
Heparin*	As prescribed

Anti-inflammatory

Aspirin	300–325mg (1 tablet) up to 4 times daily
Budesonide inhaler*	800–1200mcg (puffed) twice daily
Cetirizine	10mg (1 tablet) daily
Colchicine*	500mcg twice daily for 7 to 14 days
Corticosteroids*	Methylprednisolone, prednisone, prednisolone, dexamethasone, betamethasone, as prescribed
Cyproheptadine*	4–8mg 3 to 4 times daily, or as prescribed
Famotidine	20mg daily, or as prescribed
Ibuprofen	400 mg up to 3 times daily
Loratadine	10mg (1 tablet) daily
Montelukast*	10mg daily for 14 days, or as prescribed
Naproxen*	220mg (1 tablet) twice daily
Promethazine*	10mg twice or three times daily

Symptom support

<i>For mucus</i>	
Nebulised solutions	Various, e.g. sodium bicarb, colloidal silver nebulisation for 30 minutes 4 x daily
Bromexine	8mg three times daily
N-acetylcysteine	1 tablet (600mg) once or twice a day

Symptom support (continued)

<i>For fever</i>	
Acetaminophen (Paracetamol, Tylenol)	500mg up to four times daily
<i>For cough</i>	
Salbutamol syrup*	1 teaspoon up to three times
<i>For aches and pain</i>	
See ibuprofen and aspirin above (anti-inflammatory medicines)	

Other prescription medicines

Atorvastatin*	40mg daily
Fluvoxamine*	50mg daily or twice daily
Finasteride*	5mg daily
Fenofibrate*	160mg daily
Budesonide*	As prescribed
Decahedron*	As prescribed

Equipment

To monitor oxygen saturation	Pulse oximeter	If less than 94%, go to hospital.
To check body temperature	Thermometer	6 to 8-hourly or as needed
To relieve chest symptoms	Nebuliser	Four times daily with solutions as above
To relieve chest symptoms	Oxygen concentrator	For shortness of breath

*By prescription only in most countries.

All listed therapies are given orally unless otherwise stated.

Ivermectin is also anti-inflammatory and immune-supporting. Similarly, several of the items listed under Immune Support also have anti-inflammatory and/or antiviral properties, e.g. neem and Lugol's iodine.

Mouth and nasal hygiene solutions can be bought over-the-counter but a simple solution of sodium bicarbonate, salt and purified water can be made at home.

For the latest Covid-19 information or to learn more about early treatment options, please visit WCH member sites:

- **BiRD Group - bird-group.org/protocols**
- **FLCCC - covid19criticalcare.com/covid-19-protocols/i-mask-plus-protocol**
- **CCCA - www.canadiancovidcarealliance.org/treatment-protocols**
- **Zelenko - vladimirzelenkomd.com/treatment-protocol**
- **TFH Foundation - truthforhealth.org/patientguide**
- **Early Covid Care Experts - earlycovidcare.org**

Disclaimer:

This website and downloadable booklet are not intended to be used in place of individual medical advice. It cannot be used to diagnose illness or access treatment for web users. Individuals may use the materials provided by the World Council for Health to complement the care provided by their doctor(s). All information provided by the World Council for Health or in connection with its website is offered to promote consideration by individuals and their trained health care providers of various evidence-based prevention and treatment options. The information on this website is for general informational purposes and is not a substitute for medical advice. Errors and omissions may occur.

Stay posted on the latest guidance and preparation for children.

Subscribe to World Council for Health's Newsletter:
worldcouncilforhealth.org/subscribe



For the latest information, visit:

[WorldCouncilForHealth.org](https://www.worldcouncilforhealth.org)

Early At-Home Treatment Guide • VRS 6 • Current as of January 1, 2022